

## LECTURE 13: THE MEASUREMENT OF BAO

GONG-BO ZHAO

### 1. THE SURVEY WINDOW FUNCTION

Due to the geometry of the survey volume, we have to convolve the theoretical power spectrum with the survey window function, which is a pain, but there is no way around it.

$$(1) \quad \delta'(\mathbf{x}) = \delta(\mathbf{x})W(\mathbf{x}),$$

$$(2) \quad \tilde{\delta}'(\mathbf{k}) = \tilde{\delta}(\mathbf{k}) * \tilde{W}(\mathbf{k}),$$

where  $*$  denotes a convolution.

$$(3) \quad P'(\mathbf{k}) = \int \frac{d^3q}{(2\pi)^3} P(\mathbf{k} - \mathbf{q}) |\tilde{W}(\mathbf{q})|^2$$

Fortunately, it was found that the 3D convolution can be broken into 1D Hankel transformations, due to the convolution theorem [1].

$$(4) \quad P_\ell(k) = 4\pi(-i)^\ell \int \Delta^2 d\Delta \xi_\ell(\Delta) j_\ell(k\Delta)$$

$$(5) \quad P'_\ell(k) = 4\pi(-i)^\ell \left( \frac{2\ell+1}{2q+1} \right) \times A_{\ell,\ell'}^q \int \Delta^2 d\Delta \xi_{\ell'}(\Delta) Q_q(\Delta) j_\ell(k\Delta)$$

$$(6) \quad RR_q^{\text{tot}}(\Delta) = \frac{1}{2} \bar{n}_s^2 2\pi \Delta^3 d(\ln \Delta) Q_q(\Delta)$$

$$(7) \quad \xi'_0(\Delta) = \xi_0 Q_0 + \frac{1}{5} \xi_2 Q_2 + \frac{1}{9} \xi_4 Q_4 + \frac{1}{13} \xi_6 Q_6 + \dots$$

$$(8) \quad \begin{aligned} \xi'_2(\Delta) &= \xi_0 Q_2 + \xi_2 & \left( Q_0 + \frac{2}{7} Q_2 + \frac{2}{7} Q_4 \right) \\ &+ \xi_4 & \left( \frac{2}{7} Q_2 + \frac{100}{693} Q_4 + \frac{25}{143} Q_6 \right) \\ &+ \xi_6 & \left( \frac{25}{143} Q_4 + \frac{14}{143} Q_6 + \frac{28}{221} Q_8 \right) \end{aligned}$$




FIGURE 1. The survey window function for BOSS DR12 [6].

## 2. THE MEASUREMENTS OF BAO

**2.1. BAO using power spectrum multipoles.** The template:

$$(9) \quad \alpha_{\perp} = \frac{D_A(z)r_d^{\text{fid}}}{D_A^{\text{fid}}(z)r_d}, \quad \alpha_{\parallel} = \frac{H^{\text{fid}}(z)r_d^{\text{fid}}}{H(z)r_d}$$

$$(10) \quad P_g(k, \mu) = P_{\text{nw}}(k, \mu) \left\{ 1 + O(k)e^{-k^2[\mu^2\Sigma_{\parallel}^2 + (1-\mu^2)\Sigma_{\perp}^2]/2} \right\}$$

$$(11) \quad P_{\text{nw}}(k, \mu) = B^2 (1 + \beta\mu^2)^2 P_{\text{nw,lin}}(k) F(k, \mu)$$

$$(12) \quad F(k, \mu) = \frac{1}{(1 + k^2\mu^2\Sigma_s^2/2)}$$




FIGURE 2. The convolved power spectra [6].

$$P_\ell(k) = \left(\frac{r_s^{\text{fid}}}{r_s}\right)^3 \frac{2\ell+1}{2\alpha_\perp^2 \alpha_\parallel} \int_{-1}^1 d\mu P_g(k', \mu') \mathcal{L}_\ell(\mu) + \frac{a_{\ell 1}}{k^3} + \frac{a_{\ell 2}}{k^2} + \frac{a_{\ell 3}}{k} + a_{\ell 4} + a_{\ell 5} k$$




FIGURE 3. The BAO fit in  $k$ -space using BOSS DR12 data [6].

$$(13) \quad \begin{aligned} k' &= \frac{k(1+\epsilon)}{\alpha} \left\{ 1 + \mu^2 \left[ (1+\epsilon)^{-6} - 1 \right] \right\}^{1/2} \\ \mu' &= \frac{\mu}{(1+\epsilon)^3} \left\{ 1 + \mu^2 \left[ (1+\epsilon)^{-6} - 1 \right] \right\}^{-1/2} \end{aligned}$$


where

$$(14) \quad \alpha = \alpha_{\perp}^{2/3} \alpha_{\parallel}^{1/3}, \quad 1 + \epsilon = \left( \frac{\alpha_{\parallel}}{\alpha_{\perp}} \right)^{1/3}$$

## 2.2. BAO using correlation function multipoles.

$$(15) \quad P_{\ell}(k) = \frac{2\ell+1}{2} \int_{-1}^1 P(k, \mu) \mathcal{L}_{\ell}(\mu) d\mu$$

$$(16) \quad \xi_{\ell}(s) = \frac{i^{\ell}}{2\pi^2} \int k^2 P_{\ell}(k) j_{\ell}(ks) dk$$

FIGURE 4. The BAO fit in  $s$ -space using BOSS DR12 data [7].

$$(17) \quad \xi(s, \mu) = \sum_{\ell} \xi_{\ell}(s) \mathcal{L}_{\ell}(\mu)$$

$$(18) \quad \xi_{\ell}(s, \alpha_{\perp}, \alpha_{\parallel}) = \frac{2\ell+1}{2} \int_{-1}^1 \xi(s', \mu') \mathcal{L}_{\ell}(\mu) d\mu$$

where

$$(19) \quad s' = s \sqrt{\mu^2 \alpha_{\parallel}^2 + (1 - \mu^2) \alpha_{\perp}^2}; \quad \mu' = \mu \alpha_{\parallel} / \sqrt{\mu^2 \alpha_{\parallel}^2 + (1 - \mu^2) \alpha_{\perp}^2}$$

$$(20) \quad A_{\ell}(s) = \frac{a_{\ell,1}}{s^2} + \frac{a_{\ell,2}}{s} + a_{\ell,3}$$

$$(21) \quad \begin{aligned} \xi_0^{\text{mod}}(s) &= B_0 \xi_0(s, \alpha_{\perp}, \alpha_{\parallel}) + A_0(s) \\ \xi_2^{\text{mod}}(s) &= \xi_2(s, \alpha_{\perp}, \alpha_{\parallel}) + A_2(s) \end{aligned}$$

With this methodology, the BAO distance has been well measured from SDSS, 2dF [2, 3, 4, 5, 6, 7, 8] and DESI surveys [9, 10, 11], especially with the BAO-reconstruction technique [12, 13].

## REFERENCES

- [1] M. J. Wilson, J. A. Peacock, A. N. Taylor and S. de la Torre, “Rapid modelling of the redshift-space power spectrum multipoles for a masked density field,” *Mon. Not. Roy. Astron. Soc.* **464**, no. 3, 3121 (2017) doi:10.1093/mnras/stw2576 [arXiv:1511.07799 [astro-ph.CO]].
- [2] D. J. Eisenstein *et al.* [SDSS], “Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies,” *Astrophys. J.* **633**, 560-574 (2005) doi:10.1086/466512 [arXiv:astro-ph/0501171 [astro-ph]].
- [3] S. Cole *et al.* [2dFGRS], “The 2dF Galaxy Redshift Survey: Power-spectrum analysis of the final dataset and cosmological implications,” *Mon. Not. Roy. Astron. Soc.* **362**, 505-534 (2005) doi:10.1111/j.1365-2966.2005.09318.x [arXiv:astro-ph/0501174 [astro-ph]].
- [4] W. J. Percival *et al.* [SDSS], “Baryon Acoustic Oscillations in the Sloan Digital Sky Survey Data Release 7 Galaxy Sample,” *Mon. Not. Roy. Astron. Soc.* **401**, 2148-2168 (2010) doi:10.1111/j.1365-2966.2009.15812.x [arXiv:0907.1660 [astro-ph.CO]].
- [5] S. Alam *et al.* [BOSS], “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample,” *Mon. Not. Roy. Astron. Soc.* **470**, no. 3, 2617-2652 (2017) doi:10.1093/mnras/stx721 [arXiv:1607.03155 [astro-ph.CO]].
- [6] G. B. Zhao *et al.* [BOSS Collaboration], “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: tomographic BAO analysis of DR12 combined sample in Fourier space,” *Mon. Not. Roy. Astron. Soc.* **466**, no. 1, 762 (2017) doi:10.1093/mnras/stw3199 [arXiv:1607.03153 [astro-ph.CO]].
- [7] Y. Wang *et al.* [BOSS Collaboration], “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: tomographic BAO analysis of DR12 combined sample in configuration space,” *Mon. Not. Roy. Astron. Soc.* **469**, no. 3, 3762 (2017) doi:10.1093/mnras/stx1090 [arXiv:1607.03154 [astro-ph.CO]].
- [8] G. B. Zhao *et al.*, “The clustering of the SDSS-IV extended Baryon Oscillation Spectroscopic Survey DR14 quasar sample: a tomographic measurement of cosmic structure growth and expansion rate based on optimal redshift weights,” *Mon. Not. Roy. Astron. Soc.* **482**, no. 3, 3497 (2019) doi:10.1093/mnras/sty2845 [arXiv:1801.03043 [astro-ph.CO]].
- [9] A. G. Adame *et al.* [DESI], “DESI 2024 III: baryon acoustic oscillations from galaxies and quasars,” *JCAP* **04**, 012 (2025) doi:10.1088/1475-7516/2025/04/012 [arXiv:2404.03000 [astro-ph.CO]].
- [10] A. G. Adame *et al.* [DESI], “DESI 2024 IV: Baryon Acoustic Oscillations from the Lyman alpha forest,” *JCAP* **01**, 124 (2025) doi:10.1088/1475-7516/2025/01/124 [arXiv:2404.03001 [astro-ph.CO]].
- [11] M. Abdul Karim *et al.* [DESI], “DESI DR2 Results II: Measurements of Baryon Acoustic Oscillations and Cosmological Constraints,” [arXiv:2503.14738 [astro-ph.CO]].
- [12] D. J. Eisenstein, H. j. Seo, E. Sirkov and D. Spergel, “Improving Cosmological Distance Measurements by Reconstruction of the Baryon Acoustic Peak,” *Astrophys. J.* **664**, 675-679 (2007) doi:10.1086/518712 [arXiv:astro-ph/0604362 [astro-ph]].
- [13] E. Paillas *et al.* [DESI], “Optimal reconstruction of baryon acoustic oscillations for DESI 2024,” *JCAP* **01**, 142 (2025) doi:10.1088/1475-7516/2025/01/142 [arXiv:2404.03005 [astro-ph.CO]].